

INTRINSIC MOMENT OF INERTIA OF MEMBRANES AS BOUNDS FOR THE MASS GAP OF YANG-MILLS THEORIES.

M.P. GARCIA DEL MORAL¹, L. NAVARRO ², A. J. PÉREZ A.³ AND A. RESTUCCIA⁴

ABSTRACT. We obtain the precise condition on the potentials of Yang-Mills theories in $0+1$ dimensions and $D0$ brane quantum mechanics ensuring the discreteness of the spectrum. It is given in terms of a moment of inertia of the membrane. From it we obtain a bound for the mass gap of any $D+1$ Yang-Mills theory in the slow-mode regime. In particular we analyze the physical case $D=3$. The quantum mechanical behavior of the theories, concerning its spectrum, is determined by harmonic oscillators with frequencies given by the inertial tensor of the membrane. We find a class of quantum mechanic potential polynomials of any degree, with classical instabilities that at quantum level have purely discrete spectrum.

1. INTRODUCTION

The characterization of the discreteness of bosonic potentials in quantum mechanics represents a very challenging problem by itself and in relation to matrix model regularizations of field theories and to dimensional reduction of Yang-Mills to $0+1$ dimensions. It would provide information about the spectrum of quantum non-abelian gauge theories, as well as theories containing gravity (bosonic membrane of M-theory).

The potential of the 2-brane theories is quartic on the fields and exactly the same, after a $SU(N)$ regularization, to the dimensional reduction of Yang-Mills theories to 1 dimension (time). The potential in terms of the membrane fields has valleys extending to infinity along the directions of zero potential. There are configurations, which have been called string-like spikes [1, 2] in the context of $D=11$ supermembrane theory, for which the energy is zero. They can be attached to the membrane and even connect two membranes without any change in the energy of the system. Those configurations correspond from the Yang-Mills point of view to gauge fields valued on abelian subalgebras of $SU(N)$, in particular on the Cartan subalgebra. A toy model for these class of potentials was first studied in [3] and then extended to a toy model of the $D=11$ supersymmetric membrane in [1]. The potentials are classically unstable, quantum mechanically stable and its supersymmetric extension quantum unstable. Although the bosonic potential presents flat directions extending to infinity the walls of the valleys become narrower as we move to infinity in a way that the wave function cannot escape to infinity. A proof of discreteness of the spectrum of the $SU(N)$ regularized hamiltonian of the

Date: February 1, 2008.

bosonic membrane was presented in [4], where a bound

$$(1) \quad \langle \Psi, H\Psi \rangle \geq \langle \Psi, \lambda\Psi \rangle$$

in terms of a function $\lambda(x)$, with $\lambda(x) \rightarrow \infty$ as $\|x\| \rightarrow \infty$ was obtained. It is a sufficient condition for the discreteness of the spectrum. However in order to understand the exact property of the membrane, and, as a consequence of Yang-Mills, ensuring the discreteness of the spectrum one should look for a necessary and sufficient condition. That condition was discovered by A.M. Molchanov, [5] and recently extended by Maz'ya and Shubin [6], and makes use of the mean value, in the sense of Molchanov, of the potential on an n -dimensional cube. If the mean value goes to infinity at large distances in configuration space the spectrum is discrete, otherwise is continuous. We consider the Molchanov condition for the membrane potential. By performing the explicit calculation we obtain that at large distances in the direction of the valleys, the mean value of the potential behaves as an harmonic oscillator with frequencies determined by a moment of inertia of the membrane. There is a inertial tensor associated to the potential of the membrane, or equivalently of the $D0$ mechanics, which is strictly positive. One may interpret the mean value of the potential as a rotational energy of $D0$ branes.

This is the precise property ensuring the discreteness of the membrane and of Yang-Mills in $0 + 1$ dimensions. This approximation for Yang-Mills describe the spectrum of the theory when only the modes with zero momenta but non-zero energy are kept. This regime has been thought to contain relevant information of the IR phase of the theory in which the gluons are expected to be confined forming bound states called glueballs.

We then obtain bounds for the Yang-Mills hamiltonian in terms of the moment of inertia of the membrane. The hamiltonian bound implies that the resolvent of the Schrödinger operator is compact. It also provide a bound for the mass gap. The same results are proven for the Yang-Mills theories in the slow mode regime and give an strong indication that the complete Yang-Mills theories in D -dimensions have a mass gap.

The interesting properties of the membrane and Yang-Mills potentials are also present in more general potentials, of higher degree than four. We discuss a class of these potentials in section 3, in particular they are related to the $D = 11$ 5-brane potential. We prove discreteness of these class of potentials.

The paper is organized as follows: In section 2 some preliminary results are presented. In section 3 we obtain a class of potentials for which the spectrum of the Schrödinger operator is discrete, extending previous results. In section 4 we introduce the moment of inertia of the membrane. In section 5 we obtain a lower bound to the hamiltonian in terms of it. In section 6. the discreteness of the Yang-Mills hamiltonian in the slow mode regime is proved. In section 7 we present the conclusions.

2. PRELIMINARY RESULTS

In 1953 A. M. Molchanov (see [5]) proved that in dimension $n = 1$, the discreteness of the spectrum of the Schrödinger operator $-\Delta + V$ in $L^2(\mathbb{R}^n)$ with a locally

integrable and semi-bounded below potential V is equivalent to

$$\text{For every fixed } d > 0 \quad \int_{Q_d} V(x) dx \rightarrow \infty \quad \text{as } Q_d \rightarrow \infty$$

where Q_d is an open cube with the edges parallel to coordinate axes and of edge length d and $Q_d \rightarrow \infty$ means that the distance from Q_d to 0 goes to infinity. In the same paper Molchanov showed that this condition is not sufficient for dimension $n \geq 2$, his counter-example is based on theorem 2.1 of his paper:

The spectrum of the Schrödinger operator $-\Delta + V$ is not discret if and only if there exist a constant $C > 0$ and a sequence of disjoint cubes $\{Q_k\}_{k=1}^\infty$ with the same edge length such that for all $k = 1, 2, \dots$

$$\lambda(Q_k) = \inf \int_{Q_k} (|\nabla \psi(x)|^2 + V(x)|\psi(x)|^2) dx < C$$

where the infimum is taken over all the functions ψ such that

$$\psi|_{\partial Q_k} = 0 \text{ and } \int_{Q_k} |\psi(x)|^2 dx = 1.$$

After, Molchanov stated and proved a modified necessary and sufficient condition for the discreteness of the spectrum of Schrödinger operators for $n \geq 2$, which involve the class of negligible compact subsets of the cube. We will only state the V. Maz'ya and M. Shubin's generalization (see [6]). Before we state this theorem, let us remember the following definition of capacity. For the details and the full-general version of the theorem see [6].

Definition 1. Let $n \geq 3$, $F \subset \mathbb{R}^n$ be compact, and $Lip_c(\mathbb{R}^n)$ the set of all real-value functions with compact support satisfying a uniform Lipschitz condition in $\mathbb{R}^n$. Then the Wiener's capacity of F is defined by

$$\text{cap}(F) = \text{cap}_{\mathbb{R}^n}(F) = \inf \left\{ \int_{\mathbb{R}^n} |\nabla u(x)|^2 dx \mid u \in Lip_c(\mathbb{R}^n), u|_F = 1 \right\}$$

In physical terms the capacity of the set $F \subset \mathbb{R}^n$ is defined as the electrostatic energy over $\mathbb{R}^n$ when the electrostatic potential is set to 1 on F .

Definition 2. Let $\mathcal{G}_d \subset \mathbb{R}^n$ be an open, bounded and star-shaped set of diameter d , let $\gamma \in (0, 1)$. The *negligibility class* $\mathcal{N}_\gamma(\mathcal{G}_d; \mathbb{R}^n)$ consists of all compact sets $F \subset \overline{\mathcal{G}_d}$ satisfying $\text{cap}(F) \leq \gamma \text{cap}(\overline{\mathcal{G}_d})$.

Balls and cubes in $\mathbb{R}^n$ are useful examples of such $\mathcal{G}_d$. In what follows we denote the ball of diameter d and center x by $B_d(x)$ and the n -dimensional Lebesgue measure by $\text{Vol}(\cdot)$.

Theorem 1 (Maz'ya and Shubin). *Let $V \in L^1_{loc}(\mathbb{R}^n)$, $V \geq 0$.*

Necessity: If the spectrum of $-\Delta + V$ in $L^2(\mathbb{R}^n)$ is discrete then for every function $\gamma : (0, +\infty) \rightarrow (0, 1)$ and every $d > 0$

$$(2) \quad \inf_{F \in \mathcal{N}_\gamma(\mathcal{G}_d; \mathbb{R}^n)} \int_{\mathcal{G}_d \setminus F} V(x) dx \rightarrow +\infty \quad \text{as } \mathcal{G}_d \rightarrow \infty.$$

Sufficiency: Let a function $d \mapsto \gamma(d) \in (0, 1)$ be defined for $d > 0$ in a neighborhood of 0 and satisfying

$$\limsup_{d \downarrow 0} d^{-2} \gamma(d) = +\infty.$$

Assume that there exists $d_0 > 0$ such that (2) holds for every $d \in (0, d_0)$. Then the spectrum of $-\Delta + V$ in $L^2(\mathbb{R}^n)$ is discrete.

Remark 2. It follows from the previous theorem that a necessary condition for the discreteness of spectrum of $-\Delta + V$ is

$$(3) \quad \int_{\mathcal{G}_d} V(x) dx \rightarrow \infty \quad \text{as} \quad \mathcal{G}_d \rightarrow \infty.$$

Let's recall that in 1934, K. Friedrichs (see [6] for further references) proved that the spectrum of the Schrödinger operator $-\Delta + V$ in $L^2(\mathbb{R}^n)$ with a locally integrable potential V is discrete provided $V(x) \rightarrow \infty$ as $|x| \rightarrow \infty$.

Remark 3. The presence of negligibility sets F in the formulation of the Maz'ya and Shubin's theorem is related to the first term in the expression of $\lambda(Q_k)$. In fact this term describes also an electrostatic energy, when $\Psi = 1$ on $F \subset Q$ the energy is minimized by the harmonic solution to the corresponding Dirichlet problem. The energy of the harmonic solution defines the capacity of the capacitor $[F, Q]$ while the capacity of F is the capacity of the capacitor $[F, \infty]$.

The following lemma is very usefull tool in the next sections.

Lemma 4. For each given $\mathcal{G}_d = \mathcal{G}_d(x_0)$,

$$c_d := \inf_{F \in \mathcal{N}_\gamma(\mathcal{G}_d; \mathbb{R}^n)} \text{Vol}(\mathcal{G}_d \setminus F) > 0.$$

Proof. Let $V(x) = |x|$. Then by Friedrichs theorem the spectrum of $-\Delta + V$ is discrete, so by theorem 1 we have

$$\inf_{F \in \mathcal{N}_\gamma(\mathcal{G}_d; \mathbb{R}^n)} \int_{\mathcal{G}_d \setminus F} V(x) dx \rightarrow \infty \quad \text{as} \quad |x_0| \rightarrow \infty.$$

Now $\int_{\mathcal{G}_d \setminus F} V(x) dx \leq (|x_0| + d) \text{Vol}(\mathcal{G}_d \setminus F)$ implies that

$$\inf_{F \in \mathcal{N}_\gamma(\mathcal{G}_d; \mathbb{R}^n)} \int_{\mathcal{G}_d \setminus F} V(x) dx \leq (|x_0| + d) \inf_{F \in \mathcal{N}_\gamma(\mathcal{G}_d; \mathbb{R}^n)} \text{Vol}(\mathcal{G}_d \setminus F),$$

from which follows that $\inf_{F \in \mathcal{N}_\gamma(\mathcal{G}_d; \mathbb{R}^n)} \text{Vol}(\mathcal{G}_d \setminus F) > 0$, as we claimed. $\square$

3. SOME POTENTIALS FOR WHICH THE SPECTRUM OF ITS SCHRÖDINGER OPERATOR IS DISCRETE

In this section we use the sufficiency conditions for the discreteness of spectrum provided by theorem 1.

Theorem 5. Let $V \geq 0$, $V \in L^1_{loc}(\mathbb{R}^n)$ and c_d as in lemma 4. Suppose that there is some $d_0 > 0$ such that for all $d \in (0, d_0)$, there exists an open neighborhood Ω_d of $V^{-1}(0)$ with the following properties:

- (i) In $\mathbb{R}^n \setminus \Omega_d$, $V(x) \rightarrow \infty$ as $|x| \rightarrow \infty$.
- (ii) $\text{Vol}(B_d \cap \Omega_d) < \frac{c_d}{2}$ for every ball $B_d(x_0)$ with $|x_0| \gg d_0$.

Then the spectrum of the Schrödinger operator $-\Delta + V$ in $L^2(\mathbb{R}^n)$ is discrete.

Proof. Let $\mathcal{G}_d = B_d(x_0)$ with $d \in (0, d_0)$ fixed. Let $F \in \mathcal{N}_\gamma(\mathcal{G}_d; \mathbb{R}^n)$ and $E = (\mathcal{G}_d \setminus F) \setminus (\mathcal{G}_d \cap \Omega_d)$. Notice that $E \subset \mathbb{R}^n \setminus \Omega_d$. By condition (i) for each $M > 0$, there exists $R > 0$ such that

$$\text{If } x \in \mathbb{R}^n \setminus \mathcal{G}_d \text{ and } |x| > R, \text{ then } V(x) > M.$$

In this way, if $|x_0| - d > R$ then $|x| > R$ for all $x \in \mathcal{G}_d$, and as $V \geq 0$ we have

$$\int_{\mathcal{G}_d \setminus F} V(x) dx \geq \int_E V(x) dx \geq M \text{Vol}(E) \geq M \left(\text{Vol}(\mathcal{G}_d \setminus F) - \frac{c_d}{2} \right)$$

and by the lemma 4

$$\inf_{F \in \mathcal{N}_\gamma(\mathcal{G}_d; \mathbb{R}^n)} \int_{\mathcal{G}_d \setminus F} V(x) dx \geq M \frac{c_d}{2} \quad \text{if } |x_0| > R + d.$$

This together with theorem 1 implies that the spectrum of $-\Delta + V$ is discrete. $\square$

Now, let us apply the previous theorem.

Proposition 6. *Let $V(x) = \prod_{k=1}^n |x_k|^{\alpha_k}$, where $\alpha_k > 0$ for all $k = 1, 2, \dots, n$. Then the spectrum of the Schrödinger operator $-\Delta + V$ in $L^2(\mathbb{R}^n)$ is discrete.*

Proof. Let $d_0 = \frac{n2^n}{\omega_n}$, where ω_n denotes the area of the unit ball, and let $d \in (0, d_0)$ fixed. Consider $U_k = \{x \in \mathbb{R}^n \mid |x_k| < \epsilon_d\}$, where $0 < \epsilon_d < 1$ will be determined below. Let $\Omega_d = \cup_{k=1}^n U_k$. Then $V^{-1}(0) \subset \Omega_d$.

(a) In $\mathbb{R}^n \setminus \Omega_d$, $V(x) \rightarrow \infty$ as $|x| \rightarrow \infty$:

First, $x \in \mathbb{R}^n \setminus \Omega_d$ implies $|x_k| \geq \epsilon_d$ for all $k = 1, 2, \dots, n$.

Let $|x_j| = \max_{k=1, \dots, n} |x_k|$, $\alpha_M = \max_{k=1, \dots, n} \alpha_k$ and $\alpha_m = \min_{k=1, \dots, n} \alpha_k$. Then,

denoting $|\alpha| = \alpha_1 + \dots + \alpha_n$ we have

$$V(x) = \prod_{k=1}^n |x_k|^{\alpha_k} \geq \epsilon_d^{|\alpha| - \alpha_j} \left(\max_{k=1, \dots, n} |x_k| \right)^{\alpha_j} \geq \epsilon_d^{|\alpha| - \alpha_m} \left(\frac{|x|}{\sqrt{n}} \right)^{\alpha_j}.$$

Thus, for $|x| > 1$ we have

$$V(x) \geq \frac{\epsilon_d^{|\alpha| - \alpha_m}}{(\sqrt{n})^{\alpha_m}} |x|^{\alpha_m},$$

from which follows that $V(x) \rightarrow \infty$ as $|x| \rightarrow \infty$.

(b) $\text{Vol}(B_d \cap \Omega_d) < \frac{c_d}{2}$ for every ball $B_d(x_0)$ with $|x_0| \gg d_0$:

For $|x_0|$ large enough, if $B_d(x_0) \cap \Omega_d \neq \emptyset$ then $B_d(x_0) \cap U_k \neq \emptyset$ for only one $k \in \{1, 2, \dots, n\}$, namely $k = j$. In consequence,

$$B_d(x_0) \cap \Omega_d \subset \{x \in \mathbb{R}^n \mid |x_j| < \epsilon_d, |x_k - x_{0j}| < d, k = 1, \dots, j-1, j+1, \dots, n\},$$

from which follows that $\text{Vol}(B_d(x_0) \cap \Omega_d) \leq 2^n \epsilon_d d^{n-1}$ and this, in turn, implies what we wanted to prove if we choose

$$(4) \quad \epsilon_d < \frac{c_d}{2^n d^{n-1}}.$$

Thus, by the previous theorem we conclude that the spectrum of $-\Delta + V$ is discrete. $\square$

Remark 7. *Given that $c_d \leq \frac{\omega_n d^n}{n}$ then inequality (4) implies that $\epsilon_d < \frac{\omega_n d}{2^n n}$, which tends to zero as $d \rightarrow 0$ and also $\epsilon_d < \frac{\omega_n d}{2^n n} < \frac{\omega_n d_0}{2^n n} = \frac{\omega_n}{2^n n} \frac{n 2^n}{\omega_n} = 1$.*

4. MOMENT OF INERTIA AND DISCRETNESS OF THE POTENTIAL FOR MEMBRANE THEORIES

The matrix regularization of the bosonic membrane in the light cone gauge was done long time ago by [7, 11, 2, 8]. It consists in expanding the bosonic fields X^m in an orthonormal basis $Z_A(\sigma)$, defined on the Riemann surface Σ ,

$$(5) \quad X_m = X_m^A(\tau) Z_A(\sigma).$$

The structure constants arising from the symplectic bracket on Σ , g_{AB}^C , are those of the infinite group of area preserving diffeomorphisms.

The idea of matrix regularization is to integrate the dependence on the spatial coordinates and to regularize this model by an $SU(N)$ one, in such a way that we end up with a theory of quantum mechanics of matrices, whose structure constants f_{AB}^C converge in the large N limit to the original ones g_{AB}^C , recovering the original symmetries in the large N limit. This limit is in general not rigorously known, and a discussion on it will be presented in a separate paper [12]. In this paper we are only concerned with the theory at finite N . The potential associated to the membrane is described by

$$(6) \quad V = (Y_m^A Y_n^B f_{AB}^C)^2,$$

where Y_m , $m = 1, \dots, d$, denote the transverse direction to the light cone coordinates $Y^+ Y^-$. A, B, C denote $SU(N)$ indices and the structure constants,

$$(7) \quad f_{AB}^C = -2iN \sin\left(\frac{A \wedge B}{N} \pi\right) \delta_{A+B}^C,$$

$A = (a_1, a_2)$ $a_1, a_2 = 1, \dots, N-1$ modulo N and with $|m| + |n| \neq 0$. The structure constants in the large N limit are proportional to the structure constants of the area preserving diffeomorphisms, the residual gauge symmetry of the membranes in the light cone gauge. The large N limit is defined by taking indices A, B on a fixed set Λ and taking $N \rightarrow \infty$. We are interested in the evaluation of

$$\inf_{F \in \mathcal{N}_\gamma(\mathcal{G}_d; \mathbb{R}^n)} \int_{\mathcal{G}_d \setminus F} V,$$

where $\mathcal{G}_d$ is the star-shaped set and F the negligible set satisfying $\text{cap } F \leq \gamma \text{cap } \mathcal{G}_d$ defined in section 1, as the distance of $\mathcal{G}_d$, to a fixed point in $\mathbb{R}^n$ goes to infinity. V satisfy the assumptions of theorem 1. The integration is performed with the Lebesgue measure dY . We introduce for a given $F \in \mathcal{N}_\gamma(\mathcal{G}_d; \mathbb{R}^n)$, the center of mass of the volume $\mathcal{G}_d \setminus F$ by

$$(8) \quad \int_{\mathcal{G}_d \setminus F} Y_m^A dY = X_m^A \text{Vol}(\mathcal{G}_d \setminus F).$$

We decompose,

$$(9) \quad Y_m^A = X_m^A + x_m^A,$$

with

$$(10) \quad \int_{\mathcal{G}_d \setminus F} x_m^A dY = \int_{[\mathcal{G}_d \setminus F]_0} x_m^A dx = 0$$

where $[\mathcal{G}_d \setminus F]_0$ denote the set with its center of mass at the origin of coordinates. We then have

$$(11) \quad V(Y) = \sum_m \overline{(X_m^A + x_m^A)} D_{AB}^m (X_m^B + x_m^B),$$

where

$$(12) \quad \begin{aligned} D_{AB}^m &\equiv \sum_{n \neq m} (X_n^C + x_n^C) \overline{(X_n^D + x_n^D)} f_{BC}^E \bar{f}_{AD}^E = \\ &= 4N^2 \sum_{n \neq m} (X_n^C + x_n^C) \overline{(X_n^D + x_n^D)} \sin\left(\frac{B \wedge C}{N} \pi\right) \sin\left(\frac{A \wedge D}{N} \pi\right) \delta_{B+D}^{A+C}, \end{aligned}$$

and

$$(13) \quad V(Y) = \sum_m [(\bar{X}_m^A D_{AB}^m X_m^B + O(X_m))].$$

Here $\lim_{\|X_m\| \rightarrow \infty} \frac{O(X_m)}{X_m} = 0$ assuming the other coordinates of the center of mass bounded. We will consider first the case when only one X_m tends to infinity, namely $X_{m_0} \rightarrow \infty$, while all the others $X_m, m \neq m_0$ remain bounded. We get

$$(14) \quad \begin{aligned} \int_{\mathcal{G}_d \setminus F} V(Y) dY &= \int_{[\mathcal{G}_d \setminus F]_0} V(X + x) dx = \sum_m \left[(\bar{X}_m^A) \left(\int_{[\mathcal{G}_d \setminus F]_0} D_{AB}^m dx \right) X_m^B + O(X_m) \right] = \\ &= \sum_m \bar{X}_m^A \left[\mathbb{I}_{AB}^m + 4N^2 \sum_{n \neq m} X_n^C \bar{X}_n^D \sin\left(\frac{A \wedge C}{N} \pi\right) \sin\left(\frac{B \wedge D}{N} \pi\right) \delta_{B+D}^{A+C} \text{Vol}(\mathcal{G}_d \setminus F) \right] X_m^B \\ &\quad + O(X_m), \end{aligned}$$

where

$$(15) \quad \mathbb{I}_{AB}^m = \sum_{n \neq m} 4N^2 \int_{[\mathcal{G}_d \setminus F]_0} x_n^C \bar{x}_n^D \sin\left(\frac{A \wedge C}{N} \pi\right) \sin\left(\frac{B \wedge D}{N} \pi\right) \delta_{A+D}^{B+C} dx$$

is an intrinsic inertial tensor of the membrane associated to the distribution $[\mathcal{G}_d \setminus F]_0$. $\mathbb{I}_{AB}^m$ is a self-adjoint positive matrix. The diagonal terms are

$$(16) \quad \mathbb{I}_{AA}^m = 4N^2 \int_{[\mathcal{G}_d \setminus F]_0} \sum_{n \neq m} x_n^C \bar{x}_n^C \sin^2\left(\frac{A \wedge C}{N} \pi\right) dx.$$

Its trace is

$$(17) \quad \text{tr } \mathbb{I}^m = \sum_A \mathbb{I}_{AA}^m = 2N^4 \int_{[\mathcal{G}_d \setminus F]_0} \sum_{n \neq m} x_n^C \bar{x}_n^C dx.$$

This is the moment of inertia of the mass distribution $\mathcal{G}_d \setminus F$ with respect to the subspace generated by X_m^A . The second term in equation (14) is quartic on the center of mass coordinates. Its contribution to the integral of the potential is the value of the potential at the center of mass times the volume of the $\mathcal{G}_d \setminus F$,

$$(18) \quad (X_m^A X_n^C f_{AC}^E)^2 \text{Vol}(\mathcal{G}_d \setminus F).$$

This term becomes zero when we take the center of mass at the bottom of the valleys, in the directions where V is zero. We then conclude that along these directions where the potential is zero we obtain,

$$(19) \quad \lim_{\|X_{m_0}\| \rightarrow \infty} \left(\inf_{F \in \mathcal{N}_\gamma(\mathcal{G}_d; \mathbb{R}^n)} \int_{\mathcal{G}_d \setminus F} V dx \right) = \lim_{\|X_{m_0}\| \rightarrow \infty} \left(\inf_{F \in \mathcal{N}_\gamma(\mathcal{G}_d; \mathbb{R}^n)} \overline{X}_{m_0}^A \mathbb{I}_{AB}^{m_0} X_{m_0}^B \right) \\ = \left(\lim_{\|X_{m_0}\| \rightarrow \infty} \|X_{m_0}\|^2 \right) \inf_{F \in \mathcal{N}_\gamma(\mathcal{G}_d; \mathbb{R}^n)} \left(\frac{\overline{X}_{m_0}^A}{\|X_{m_0}\|} \mathbb{I}_{AB}^{m_0} \frac{X_{m_0}^B}{\|X_{m_0}\|} \right),$$

and we notice that

$$\inf_{F \in \mathcal{N}_\gamma(\mathcal{G}_d; \mathbb{R}^n)} \left(\frac{\overline{X}_{m_0}^A}{\|X_{m_0}\|} \mathbb{I}_{AB}^{m_0} \frac{X_{m_0}^B}{\|X_{m_0}\|} \right) \geq \inf_{F \in \mathcal{N}_\gamma(\mathcal{G}_d; \mathbb{R}^n)} \lambda_F,$$

where λ_F is the minimum eigenvalue of $\mathbb{I}_{AB}^{m_0}$. It may be shown by explicit calculations and using lemma 4 that the infimum of the right hand side of last inequality is greater than 0. One may also obtain that conclusion in an indirect way from theorem 1. In fact we know from an independent argument (see next section or [4]), that the spectrum of the Schrödinger operator we are considering is discrete with finite multiplicity. According to theorem 1, the left hand member of equation (19) must then go to infinity when $\|X_{m_0}\| \rightarrow \infty$, hence it is necessary that

$$\inf_{F \in \mathcal{N}_\gamma(\mathcal{G}_d; \mathbb{R}^n)} \lambda_F > 0.$$

If the center of mass goes to infinity in more than one direction X_m , $m = 1, \dots, d$, (19) remains valid. The inertial tensor has now an additional index $\mathbb{I}_{AB}^{mn}$ where $m, n = 1, \dots, d$. And it is a strictly positive self adjoint matrix, the diagonal terms have the same expression (16). The corresponding equality (19) holds when the center of mass of $\mathcal{G}_d \setminus F$ is located on the directions where the potential is zero. In other directions where the potential does not vanish the left hand side of (19) is greater or equal to the quadratic right hand side of (19).

We remark that this relation is the most precise one can obtain for the quantum Schrödinger operator, since the condition of the theorem in section 1 is a necessary and sufficient condition. The mean value, in the sense of Molchanov, of the membrane potential is bounded by the mean value of an harmonic oscillator at large distances in the configuration space and it is equal to it at the directions of zero potential. This bound implies, using Maz'ya and Shubin's theorem, that the corresponding Schrödinger operator is discrete with finite multiplicity.

5. BOUNDS FOR THE HAMILTONIAN

We present in this section a bound for the hamiltonian of the regularized bosonic M2 brane in terms of the integrand of the moment of inertia we discussed in the previous section. It is an independent argument showing the discreteness of the spectrum and providing bounds to all the eigenvalues of the membrane. It represents a generalization of the proof given in [3].

Hereafter we use the notation x_m^A instead of Y_m^A for the configuration variables. We re-arrange the potential of the M2 brane in the form

$$(20) \quad V(x) = \sum_m^d x_m^A \bar{x}_m^B M_{AB}^m$$

where

$$(21) \quad M_{AB}^m = \sum_{n \neq m} f_{AE}^C x_n^E \bar{f}_{BD}^C \bar{x}_n^D$$

is a positive hermitian matrix. Then, the hamiltonian can be written as,

$$(22) \quad H = -\frac{1}{2}\Delta + \frac{1}{2}(-\Delta + 2V),$$

$$-\Delta + 2V = \sum_m \left[-\left(\frac{\partial}{\partial x_m^A} \right)^2 + 2x_m^A \bar{x}_m^B M_{AB}^m \right] \geq \sum \sqrt{2}(\text{tr } M^m)^{1/2},$$

hence

$$(23) \quad H \geq \frac{1}{2} \left(-\Delta + \sqrt{2} \sum_m (\text{tr } M^m)^{1/2} \right).$$

The computation for $\text{tr } M^m$ gives

$$(24) \quad \text{tr } M^m = 2N^4 \sum_{n \neq m} x_n^E \bar{x}_n^E.$$

We finally get the operatorial bound,

$$(25) \quad H \geq \frac{1}{2} \left(-\Delta + 2N^2 \sum_m \left(\sum_{n \neq m} (x_n^E \bar{x}_n^E)^{1/2} \right) \right) \equiv \tilde{H}.$$

The potential $\tilde{V} \equiv 2N^2 \sum_m (\sum_{n \neq m} x_n^E \bar{x}_n^E)^{1/2}$, satisfies the criteria: $\tilde{V} \rightarrow \infty$ when $\|x\| = (\sum_n x_n^E x_n^E)^{1/2} \rightarrow \infty$ implying that the resolvent of H is compact, H has a discrete spectrum with finite multiplicity. The formula also provides a bound for the mass gap of H given by the first eigenvalue of the operator $\tilde{H}$.

6. SPECTRUM OF YANG-MILLS THEORIES IN THE SLOW-MODE REGIME

In this section we will consider the explicit relation between dimensional reduced Yang-Mills theories to $0+1$ dimensions and the membrane moment of inertia condition ensuring the discreteness of the spectrum.

The hamiltonian of Yang-Mills in $D+1$ space-time dimensions has the following hamiltonian

$$(26) \quad H_{ym} = \int_{\Sigma_2} \frac{g^2}{2} (\Pi^i)^2 + \frac{1}{4g^2} F_{ij}^a F_{ij}^a - \mathcal{D}_i \Pi^{ai} A_0^{ai}$$

where Π^{ai} represents the conjugate momenta to A_i^a . The indices of color are $a = 1, \dots, \dim SU(N) = N^2 - 1$, and $i = 1, \dots, D$ labels the spatial components of gauge fields. The last piece of the Hamiltonian represents the Gauss law:

$$(27) \quad \mathcal{D}_i \Pi^{ai} = 0.$$

We assume we are in the slow mode regime and all of the fast modes has been integrated out in such a way that we deal with an effective hamiltonian containing gauge fields $A_i(t)$ without spatial dependence. This approximation has been considered to keep relevant information of the theory in the IR limit where the gluons are supposed to be confined forming glueballs bound states. The hamiltonian has then the expression,

$$(28) \quad H_{ym}^{red} = \text{tr} \left[\frac{g^2 V_m}{2} (\Pi^i)^2 + \frac{V_m}{4g^2} (f^{abc} A_i^b A_j^c)^2 - [A_i, \Pi^i]^a A_0^{ia} \right] (t).$$

In order to make comparison with the hamiltonian of the preceding sections we redefine the fields in terms of dimensionless ones. Following [10] we take,

$$(29) \quad A_i \rightarrow \frac{g^{1/3}}{V_m^{1/3}} A_i, \quad P^i \rightarrow \frac{1}{g^{2/3} V_i^{2/3}} P_i.$$

The hamiltonian can be re-written as,

$$(30) \quad H_{ym}^{red} = \frac{g^{2/3}}{V_m^{1/3}} \text{tr} \left[\frac{1}{2} (\Pi^i)^2 + \frac{1}{4} (f^{abc} A_i^b A_j^c)^2 - [A_i, \Pi^i]^a A_0^{ia} \right] (t) = \frac{g^{2/3}}{V_m^{1/3}} H.$$

The mass operator of the regularized bosonic membrane also in terms of dimensionless variables can be written as,

$$(31) \quad \frac{M^2}{2} = T^{2/3} \text{tr} \left[\frac{1}{2} (\Pi^i)^2 + \frac{1}{4} (f^{abc} A_i^b A_j^c)^2 - [A_i, \Pi^i]^a A_0^{ia} \right] (t) = T^{2/3} H.$$

This means that the two theories are equivalent and the discreteness of Yang-Mills in the slow mode regime is related to the existence of a tensor of moment of inertia of the membrane. We interpret it as an strong evidence that the configuration space for Yang-Mills theories in the large N limit is compact. The discreteness condition then is realized as a condition of "rotational energy" of $D0$ branes in the configuration space of the bosonic membrane.

6.1. Mass Gap and analytic eigenvalues distribution. An interesting problem is to estimate the mass gap of the theory for a particular value of N . We have from the previous section

$$(32) \quad H \geq \frac{1}{2} \left(-\Delta + 2N^2 \sum_m \left(\sum_{n \neq m} (x_n^E \bar{x}_n^E)^{1/2} \right) \right) \equiv \tilde{H}.$$

Let us now consider, as an example, the case of $SU(3)$ in $D = 3 + 1$ since it is the gauge group realized in the nature. The number of degrees of freedom once the gauge fixing condition has been taken into account is $d = 2$, $A = 1, \dots, 8$, and $N = 3$. $\tilde{H}$ is bounded from below by the following hamiltonian

$$(33) \quad H_1 = -\Delta + 2N^2 (X^I X^I)^{1/2} \quad I, J = 1, \dots, 16$$

The eigenfunctions of H_1 are expressed in terms of Bessel functions. The eigenvalues of H_1 are lower bounds, one by one, of the eigenvalues of H . That is, $SU(3)$ Yang-Mills in the slow-mode regime in $3 + 1$ dimensions has a discrete spectrum and its eigenvalues and mass gap are bounded by those of $\frac{g^{2/3}}{V_m^{1/3}} H_1$.

7. CONCLUSIONS

We showed that the most precise condition ensuring the discreteness of the spectrum of the membrane theories and Yang-Mills in the slow mode regime, is given in terms of an intrinsic moment of inertia of the membrane. It may be interpreted as if the membrane, or equivalently the $D0$ branes describing it, have a rotational energy. It is a quantum mechanical effect. The condition is obtained from the Molchanov, Maz'ya and Shubin necessary and sufficient condition on the potential of a Schrödinger operator to have a discrete spectrum. The criteria is expressed not in terms of the behaviour of the potentials at each point, but by a mean value, on the configuration space. The mean value in the sense of Molchanov considers the integral of the potential on a finite region of configuration space. It can be naturally associated to a discretization of configuration space in the quantum theory. We found that the mean value in the direction of the valleys where the potential is zero, at large distances in the configuration space, is the same as a harmonic oscillator with frequencies given by the tensor of inertia of the membrane. The interesting feature is that all previously known bounds for the membrane and Yang-Mills potential were linear on the configuration variables, while our bound is quadratic on the configuration variables. The bounds we obtained should also be relevant on the analysis of the spectrum of wrapped supermembrane with nontrivial central charges ([13],[14],[15],[16]).

We also obtained bounds, based on the moment of inertia for the hamiltonian of Yang-Mills theories in the slow mode regime which allows to obtain interpretation about the mass gap of the theory. In particular a bound for $SU(3)$ in $3+1$ dimensions is given by a hamiltonian whose spectrum and eigenvalues are known, and its eigenfunctions are expressed in terms of Bessel functions.

In section 3 we obtained a class of potentials, of any degree in the configuration coordinates, presenting the same valley properties of the quartic potentials discussed in section 4, 5 and 6. In particular, one of them has similarities with the potential for the 5-brane in $D = 11$. We proved that the associated Schrödinger operator has discrete spectrum, with finite multiplicity.

ACKNOWLEDGMENTS

We are very grateful to L. Boulton, F. Cachazo, H. Nicolai, Sh. Matsuura, V. Strauss for helpful discussions and comments; to C. Burgess, F. Cachazo, R. Myers, F. Quevedo, for their great support and encouragement. M.P.G.M. and A.R. would like to thank Perimeter Institute for kind hospitality while part of this work was done. M.P.G.M. acknowledge support from NSERC of Canada and MEDT of Ontario, and NSERC Discovery grant.

REFERENCES

- [1] B. de Wit , M. Luscher , H. Nicolai *The Supermembrane Is Unstable*. Nucl.Phys.**B320**:135,1989
- [2] H Nicolai, R Helling *Supermembranes and M(atr)ix theory*. [hep-th/9809103](#)
- [3] B. Simon, Ann. Phys 146 (1983)209
- [4] M. Lüscher Some Analytic Results Concerning The Mass Spectrum Of Yang-Mills Gauge Theories On A Torus. Nucl.Phys.**B219**:233-261,1983
- [5] A.M.Molchanov *On the discreteness of the spectrum conditions for selfadjoint differential equations of the second order*, Proc. Moscow Math. Society **2** (1953) 169-199 (Russian).
- [6] Mazy'a, V. and Shubin, M. Discreteness of spectrum and positivity criteria for Schrödinger operators. Annals of Mathematics, **162** (2005), 919-942.

- [7] B. de Wit, J. Hoppe, H. Nicolai *On The Quantum Mechanics Of Supermembranes* Nucl.Phys. **B305**:545,1988.
- [8] M. Claudson, M. B. Halpern *Supersymmetric Ground State Wave Functions.* Nucl.Phys.**B250**:689,1985
- [9] B. de Wit, Marquard and H. Nicolai, *Area Preserving Diffeomorphisms And Supermembrane Lorentz Invariance* Commun.Math.Phys. **128** (1990)39-62.
- [10] G. Gabadadze *Modeling The Glueball Spectrum By A Closed Bosonic Membrane* Phys.Rev.**D58**(1998)094015 [hep-ph/9710402](#)
- [11] J. Hoppe, MIT Ph.D Thesis, 1982.
- [12] L.Boulton, M.P.Garcia del Moral and A. Restuccia. In preparation.
- [13] J. Bellorin, A. Restuccia [hep-th/0510259](#), *D=11 Supermembrane wrapped on calibrated sub-manifolds* Nucl.Phys. **B737** (2006) 190-208
- [14] L. Boulton, M.P. Garcia del Moral, A. Restuccia *Discreteness of the spectrum of the compactified D=11 supermembrane with non-trivial winding* Nucl.Phys. **B671** (2003) 343-358, [hep-th/0211047](#)
- [15] L. S. Boulton, M. P. Garcia del Moral, I. Martin, A. Restuccia *On the spectrum of a matrix model for the D=11 supermembrane compactified on a torus with non-trivial winding* Class.Quant.Grav. **19** (2002) 2951 [hep-th/0109153](#)
- [16] M.P.Garcia del Moral, A. Restuccia *On The spectrum of a Noncommutative Formulation of the D=11 Supermembrane with Winding* Phys.Rev. D66 (2002) 045023 [hep-th/0103261](#)

¹ PERIMETER INSTITUTE FOR THEORETICAL PHYSICS, WATERLOO, CANADA, ONTARIO N@J 2Y5, CANADA, DEPT. OF PHYSICS AND ASTRONOMY, MACMASTER UNIVERSITY, 1280 MAIN STREET WEST, HAMILTON, ONTARIO, L8S 4M1, CANADA, DAMTP, DAMTP, CENTRE FOR MATHEMATICAL SCIENCES, UNIVERSITY OF CAMBRIDGE CB3 0WA, UK.

E-mail address: mmoral@perimeterinstitute.ca, M.G.d.Moral@damtp.cam.ac.uk

² DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD SIMÓN BOLÍVAR, APARTADO 89000, CARACAS 1080-A, VENEZUELA.

E-mail address: lnavarro@ma.usb.ve

³ DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD SIMÓN BOLÍVAR, APARTADO 89000, CARACAS 1080-A, VENEZUELA.

E-mail address: ajperez@ma.usb.ve

⁴ DEPARTAMENTO DE FÍSICA, UNIVERSIDAD SIMÓN BOLÍVAR, APARTADO 89000, CARACAS 1080-A, VENEZUELA.

E-mail address: arestu@usb.ve